List of tables

Table No.	Titles	Page No.
Table 1	Frequency and percentage distribution of baseline information	57
	of ASHA	
Table 2	Knowledge and practices level of ASHA after HBNC training	58
Table 3	Domain wise knowledge gain of ASHA on HBNC post re-	59
	education	
Table 4	Domain wise practice improvement of ASHA on HBNC post re-	62
	education	
Table 5	Improvement of knowledge, attitude and practices scores of	64
	ASHA after HBNC re-education	
Table 6	The pair wise comparison of knowledge of ASHA before and	65
	after implementation of HBNC re-education using Post Hoc	
	Bonferroni test	
Table 7	The pair wise comparison of attitude of ASHA before and after	66
	implementation of HBNC re-education using Post Hoc	
	Bonferroni test	
Table 8	The pair wise comparison of practices of ASHA before and after	66
	implementation of HBNC re-education using Post Hoc	
	Bonferroni test	
Table 9	Attitude of ASHA on Identification of high risk newborn and	68
	thermoregulation	

Table 10	Attitude of ASHA on Breastfeeding care practices	69
Table 11	Attitude of ASHA on Infection control practices	70
Table 12	Association between levels of knowledge with the selected	73
	baseline data of ASHA	
Table 13	Association between levels of attitude with the selected baseline	74
	data of ASHA	
Table 14	Association between levels of practice with the selected baseline	75
	data of ASHA	
Table 15	Correlation between knowledge, attitude and practices of ASHA	76
Table 16	Frequency and percentage distribution profile of mother and	78
	baby	
Table 17	Response of mother regarding newborn care practices	82
Table 18	Unsafe newborn care practices of mothers	86
Table 19	Response of mother regarding HBNC provided by ASHA	87

List of figures

Legends	Page No
Conceptual framework adapted from Health Belief Model	13
(Rosenstock, 1974)	
Study site – Doiwala, Dehradun, Uttarakhand	29
Schematic representation of sampling of ASHA in phase I	30
Data collection procedure for phase II	41
Schedule for data collection of the study	42
Comparison of pretest, posttest 1, posttest 2 and posttest 3 mean	60
percentage of knowledge and practice of ASHA workers.	
Comparison of pre-test and post-test mean knowledge scores of	61
ASHA	
Comparison of pre-test and post-test mean practice scores of	63
ASHA	
Comparison of pre-test and post-test mean attitude scores of	64
ASHA	
Correlation of knowledge and attitude of ASHA	76
Correlation of knowledge and practice of ASHA	77
Correlation of practice and attitude of ASHA	77
The most unsafe newborn care practices of the mothers (in %)	86
	Conceptual framework adapted from Health Belief Model (Rosenstock, 1974) Study site – Doiwala, Dehradun, Uttarakhand Schematic representation of sampling of ASHA in phase I Data collection procedure for phase II Schedule for data collection of the study Comparison of pretest, posttest 1, posttest 2 and posttest 3 mean percentage of knowledge and practice of ASHA workers. Comparison of pre-test and post-test mean knowledge scores of ASHA Comparison of pre-test and post-test mean practice scores of ASHA Comparison of pre-test and post-test mean attitude scores of ASHA Correlation of knowledge and attitude of ASHA Correlation of knowledge and practice of ASHA Correlation of practice and attitude of ASHA